

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b>  <b>A61K 7/48</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/10755</b> <b>(43) International Publication Date:</b> 10 June 1993 (10.06.93)
<b>(21) International Application Number:</b> PCT/US92/09737 <b>(22) International Filing Date:</b> 9 November 1992 (09.11.92)  <b>(30) Priority data:</b> 796,749 25 November 1991 (25.11.91) US  <b>(71) Applicant:</b> RICHARDSON-VICKS, INC. [US/US]; One Far Mill Crossing, Shelton, CT 06484 (US).  <b>(72) Inventors:</b> BLANK, Roy, Lonnie ; 39 Pennington Way, Spring Valley, NY 10977 (US). DOUGHTY, Darell, Gene ; 11841 Nathanshill Lane, Cincinnati, OH 45249 (US). LINARES, Carlos, Gabriel ; 6340 Trail Ridge Court, Loveland, OH 45140 (US).	<b>(74) Agents:</b> REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45202 (US).  <b>(81) Designated States:</b> AU, BB, BG, BR, CA, CS, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, PL, RO, RU, SD, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>	
<b>(54) Title:</b> COMPOSITIONS FOR REGULATING SKIN WRINKLES AND/OR SKIN ATROPHY		
<b>(57) Abstract</b>  The present invention relates to a composition for regulating wrinkles and/or atrophy in mammalian skin comprising treating the skin with a safe and effective amount of salicylic acid and/or additional active component.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

COMPOSITIONS FOR REGULATING  
SKIN WRINKLES AND/OR SKIN ATROPHY

5

Technical Field

The present invention relates to the field of anti-aging of skin. Specifically, the invention relates to novel compositions for effacing and preventing wrinkles in mammalian skin.

10

Background of the Invention

Skin is subject to abuse by many extrinsic (environmental) factors as well as intrinsic (chronoaging) factors. A common extrinsic factor is exposure to ultraviolet radiation. Whether extrinsic or intrinsic, the abuse results in wrinkling of the skin. To many people, skin wrinkles are a reminder of the disappearance of youth. As a result, the elimination of wrinkles has become a booming business in youth-conscious societies. Treatments range from cosmetic creams and moisturizers to various forms of cosmetic surgery.

20

Chronoaging results in the thinning and general degradation of skin. As the skin naturally ages, there is a reduction in the cells and blood vessels that supply the skin. There is also a flattening of the dermal-epidermal junction which results in weaker mechanical resistance of this junction. As a consequence, older persons are more susceptible to blister formation in cases of mechanical trauma or disease processes. (See Oikarinen, (1990) "The Aging of Skin: Chronoaging Versus Photoaging", Photodermatal. Photoimmunol. Photomed., Vol. 7, pp 3-4).

30

It is known to use salicylic acid for the treatment of acne, see for example, U.S Patents 4,891,227 and 4,891,228, to Thaman et al., both issued January 2, 1990 the disclosures of which are incorporated herein. Further, salicylic acid has been used for the removal of wart, corns and calluses; for the treatment of psoriasis, seborrheic dermatitis and dandruff; and for the topical treatment of ringworm infection. A listing of commercially available products containing salicylic acid will be found in the Physician's Desk Reference, 45th Edition, 1991, page 323. However, these prior art

35

- 2 -

uses of salicylic acid have generally involved short term treatments in which relatively large doses of the acid are applied (i.e., sufficient to cause significant irritation and often peeling) in order to obtain a cure or treatment of the particular condition, such as removal of comedones, as opposed to persistent treatment of normal aging skin.

#### Objects of the Present Invention

It is an object of the present invention to provide compositions for regulating wrinkles and/or atrophy in mammalian skin which comprise a safe and effective amount of an anti-wrinkle/anti-atrophy agent.

#### Summary of the Invention

The present invention relates to compositions for regulating wrinkles and/or atrophy in mammalian skin comprising a safe and effective amount of salicylic acid along with another active agent selected from the group consisting of a safe and effective amount of a sunscreen, an anti-inflammatory agent, a vitamin, an anti-oxidant, a chelator, a retinoid, a benzofuran derivative, an N-acetyl-L-cysteine derivative and a skin protectant, derivatives thereof and mixtures thereof.

All percentages and ratios used herein are by weight and all measurements are at 25°C unless otherwise indicated.

#### Detailed Description of the Invention

As used herein, "alkyl" means an unsubstituted carbon-containing chain which may be straight, branched or cyclic, preferably straight or branched, more preferably straight; saturated, monounsaturated (i.e., one double or triple bond in the chain), or polyunsaturated (i.e., two or more double bonds in the chain; two or more triple bonds in the chain; one or more double and one or more triple bonds in the chain), preferably saturated.

As used herein, "topical application" means directly laying on or spreading on outer skin.

As used herein, "pharmaceutically-acceptable" means that drugs, medicaments or inert ingredients which the term describes are suitable for use in contact with the tissues of humans and lower

- 3 -

animals without undue toxicity, incompatibility, instability, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio.

5 As used herein, "regulating wrinkles" means preventing, retarding, arresting, or reversing the process of wrinkle formation in mammalian skin.

10 As used herein, "skin atrophy" means the thinning and/or general degradation of the dermis often characterized by a decrease in collagen and/or elastin as well as decreased number, size and doubling potential of fibroblast cells. Skin atrophy is a natural result of aging. Skin atrophy is often an undesirable side effect resulting from treatment with corticosteroids.

15 As used herein, "regulating skin atrophy" means preventing, retarding, arresting, or reversing the process of atrophy in mammalian skin.

20 As used herein, "safe and effective amount" means an amount of compound or composition sufficient to significantly induce a positive modification in the condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgment. The safe and effective amount of the compound or composition will vary with the particular condition being treated, the age and physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, the specific  
25 compound or composition employed, the particular pharmaceutically-acceptable carrier utilized, and like factors within the knowledge and expertise of the attending physician.

30 As used herein, "chronic treatment" means continued treatment with an active agent over an extended period during a subject's lifetime, preferably for at least about three weeks, more preferably from about three months to about twenty years, more preferably from about six months to about ten years, more preferably still from about one year to about five years.

35 As used herein, all percentages are by weight unless otherwise specified.

- 4 -

Salicylic Acid Active Compound

The salicylic acid active component can be salicylic acid alone, salicylic acid derivatives and salicylic acid in combination with other active ingredients described below. Most preferred is  
5 salicylic acid in a hydroalcoholic solution.

Salicylic acid is a well known active component and is generally described in U.S. Patent 4,514,385, to Damani, et al., assigned to Alcon Laboratories, issued April 30, 1985.

The preferred topical carrier for the salicylic acid active  
10 comprises a hydroalcoholic solution at pH 2 to 4 of salicylic acid as the active anti-acne ingredient together with a specific anionic surfactant component. More preferably such active is a stable, hydroalcoholic composition having a pH value of from 2 to 4 and containing from about 0.2 to about 5.0 percent by weight of  
15 salicylic acid and from about 0.2 to about 5.0 percent by weight of sodium methyl cocoyl taurate and/or sodium methyl oleoyl taurate as the anionic surfactant component. Generally, a sufficient amount of a cosmetically acceptable alkaline component (i.e., alkalizing agent) to provide and maintain the composition with a pH from about  
20 2.0 to about 4 is included.

As the alcohol component of the hydroalcoholic solvent, from about 10 to about 60 percent by weight of ethyl alcohol, measured as total  $C_2H_5OH$  content, is preferred although a like amount of isopropyl alcohol ( $C_3H_7OH$ ) may also be beneficially utilized. From  
25 about 30 to about 80 percent by weight of water is also required as the aqueous component of the hydroalcoholic solvent.

The anionic surfactant component of this active composition, i.e., the taurate surfactant component, is specifically directed to sodium methyl cocoyl taurate and sodium methyl oleoyl taurate, both  
30 of which are readily available from diverse commercial suppliers, as noted in The Cosmetic, Toiletry and Fragrance Association (CTFA) Cosmetic Ingredient Dictionary, 3rd Edition, 1982, pages 286-287.

Although it is preferred to use the taurate surfactant as the sole surfactant in the active compositions, other surfactants may be  
35 included, the nonionic type having preference over the anionic type

- 5 -

in view of the relative non-irritating characteristic to the skin of the former. Cationic type surfactants, which are most irritating to the skin, are less preferred because of their marked susceptibility to hydrolysis at the low acidic pH of the subject compositions.

5       The pH value of the preferred active component, from about 2 to about 3.5, may be achieved by use of appropriate cosmetically acceptable primary or dual buffer systems. In most instances, the resultant pH of the hydroalcoholic solution of salicylic acid is slightly below or at the lower end of the indicated range, and all  
10       that is required to adjust the pH to a desired higher value within the indicated range is to add an alkaline additive such as is commonly utilized in cosmetic formulations for such purpose. Although sodium carbonate is preferred, other suitable alkalinizing agents include potassium carbonate, sodium hydroxide, potassium  
15       hydroxide, triethanolamine and the like. If deemed necessary to change or adjust the pH to a lower value, a suitable cosmetically acceptable acidifying agent such as citric acid may be employed.

#### Combination Actives

##### A.   Sunscreens and Sunblocks

20       Optimum regulation of skin wrinkling resulting from exposure to U.V. light can be obtained by using a combination of the salicylic acid active of the present invention together with sunscreens or sunblocks. Useful sunblocks include, for example, zinc oxide and titanium dioxide.

25       Photodamage is a predominant cause of skin wrinkling. Thus, for purposes of wrinkle prevention, the combination of the active compound with a UVA and/or UVB sunscreen would be most desirable. The inclusion of sunscreens in compositions of the present invention will provide immediate protection against acute UV damage. Thus,  
30       the sunscreen will prevent further wrinkle formation caused by UV radiation, while the active compound regulates existing wrinkles and skin atrophy.

35       A wide variety of conventional suncreening agents are suitable for use in combination with the active compound. Segarin, et al., at Chapter VIII, pages 189 et seq., of Cosmetics Science and



- 6 -

Technology, disclose numerous suitable agents. Specific suitable sunscreens include, for example: p-aminobenzoic acid, its salts and its derivatives (ethyl, isobutyl, glyceryl esters; p-dimethylaminobenzoic acid); anthranilates (i.e., o-aminobenzoates; methyl, menthyl, phenyl, benzyl, phenylethyl, linalyl, terpinyl, and cyclohexenyl esters); salicylates (amyl, phenyl, benzyl, menthyl, glyceryl, and dipropylene glycol esters); cinnamic acid derivatives (methyl and benzyl esters,  $\alpha$ -phenyl cinnamitrile; butyl cinnamoyl pyruvate); Dihydroxycinnamic acid derivatives (umbelliferone, methylumbelliferone, methylaceto-umbelliferone); trihydroxycinnamic acid derivatives (esculetin, methyl esculetin, daphnetin, and the glucosides, esculin and daphnin); hydrocarbons (diphenylbutadiene, stilbene); dibenzalacetone and benzalacetophenone; Naphthol-sulfonates (sodium salts of 2-naphthol-3,6-disulfonic and of 2-naphthol-6,8-disulfonic acids); Dihydroxy-naphthoic acid and its salts; o- and p-Hydroxybiphenyldisulfonates; Coumarin derivatives (7-hydroxy, 7-methyl, 3-phenyl); Diazoles (2-acetyl-3-bromoindazole, phenyl benzoxazole, methyl naphthoxazole, various aryl benzothiazoles); Quinine salts (bisulfate, sulfate, chloride, oleate, and tannate); Quinoline derivatives (8-hydroxyquinoline salts, 2-phenylquinoline); Hydroxy- or methoxy-substituted benzophenones; Uric and vilouric acids; Tannic acid and its derivatives (e.g., hexaethylether); (Butyl carbotoI) (6-propyl piperonyl) ether; Hydroquinone; Benzophenones (Oxybenzene, Sulisobenzene, Dioxymethane, Benzoresorcinol, 2,2',4,4'-Tetrahydroxybenzophenone, 2,2'-Dihydroxy-4,4'-dimethoxybenzophenone, Octabenzene; 4-Isopropylidibenzoylmethane; Butylmethoxydibenzoylmethane; Etocrylene; and 4-isopropyl-di-benzoylmethane.

Preferred sunscreens useful in the compositions of the present invention are 2-ethylhexyl-p-methoxycinnamate, butylmethoxydibenzoylmethane, 2-hydroxy-4-methoxybenzophenone, octyl-dimethyl-p-aminobenzoic acid and mixtures thereof.

A safe and effective amount of sunscreen may be used in the compositions of the present invention. The sunscreens must be compatible with the active compound. Generally the composition

- 7 -

may comprise from about 1% to about 20%, preferably from about 2% to about 10%, of a suncreening agent. Exact amounts will vary depending upon the sunscreen chosen and the desired Sun Protection Factor (SPF).

5 Also particularly useful in the present invention are sunscreens such as those disclosed in Sabatelli, U.S. Patent Application Serial No. 054,085\* (filed June 2, 1987) and Sabatelli et al., U.S. Patent Application Serial No 054,046\*\* (filed June 2, 1987).  
10 The suncreening agents disclosed therein have, in a single molecule, two distinct chromophore moieties which exhibit different ultra-violet radiation absorption spectra. One of the chromophore moieties absorbs predominantly in the UVB radiation range and the other absorbs strongly in the UVA radiation range.

15 An agent may also be added to any of the compositions of the present invention to improve the skin substantivity of those compositions, particularly to enhance their resistance to being washed off by water, or rubbed off. A preferred agent which will provide this benefit is a copolymer of ethylene and acrylic acid. Compositions comprising this copolymer are disclosed in U.S. Patent  
20 4,663,157, Brock, issued May 5, 1987, which is incorporated herein by reference.

#### B. Anti-Inflammatory Agents

In a preferred wrinkle and atrophy regulating composition of the present invention, an anti-inflammatory agent is included as an  
25 active agent along with the active compound. The inclusion of an anti-inflammatory agent enhances the wrinkle regulating benefits of the compositions. The anti-inflammatory agent protects strongly in the UVA radiation range (though it also provides some UVB protection as well) thereby preventing further wrinkle formation caused by UV  
30 radiation, while the active compound regulates existing wrinkles and skin atrophy. Thus the combination provides broad protection. The topical use of anti-inflammatory agents reduces photo-aging of the skin resulting from chronic exposure to UV radiation. (See U.S. Patent 4,847,071, Bissett, Bush, and Chatterjee, issued July 11, 1989, incorporated herein by reference; and U.S. Patent 4,847,069,  
35 (Equivalents = \*US Patent 4,937,370; \*\*US Patent 4,999,186.)

- 8 -

Bissett and Chatterjee, issued July 11, 1989, incorporated herein by reference.)

5 A safe and effective amount of an anti-inflammatory agent of the present invention, preferably from about 0.1% to about 10%, more preferably from about 0.5% to about 5%, of the composition. The exact amount of anti-inflammatory agent to be used in the compositions will depend on the particular anti-inflammatory agent utilized since such agents vary widely in potency.

10 Steroidal anti-inflammatory agents, including but not limited to, corticosteroids such as hydrocortisone, hydroxyltriamcinolone, alpha-methyl dexamethasone, dexamethasone-phosphate, beclomethasone dipropionate, clobetasol valerate, desonide, desoxymethasone, desoxycorticosterone acetate, dexamethasone, dichlorisone, diflorasone diacetate, diflucortolone valerate, fluadrenolone, 15 flucilorolone acetonide, fludrocortisone, flumethasone pivalate, fluosinolone acetonide, fluocinonide, flucortine butylester, flucortolone, fluprednidene (fluprednylidene) acetate, fluran-drenolone, halcinonide, hydrocortisone acetate, hydrocortisone butyrate, methylprednisolone, triamcinolone acetonide, cortisone, 20 cortodoxone, flucetonide, fludrocortisone, difluorosone diacetate, fluradrenolone acetonide, medrysone, amcinafel, amcinafide, betamethasone and the balance of its esters, chloroprednisone, chlorprednisone acetate, clocortelone, clescicolone, dichlorisone, difluprednate, fluciloronide, flunisolid, fluoromethalone, flu-perolone, fluprednisolone, hydrocortisone valerate, hydrocortisone 25 cyclopentylpropionate, hydrocortamate, meprednisone, paramethasone, prednisolone, prednisone, beclomethasone dipropionate, tri-amcinolone, and mixtures thereof may be used. The preferred steroidal anti-inflammatory for use in the present invention is 30 hydrocortisone.

A second class of anti-inflammatory agents which is useful in the compositions of the present invention includes the nonsteroidal anti-inflammatory agents. The variety of compounds encompassed by this group are well-known to those skilled in the art. For detailed 35 disclosure of the chemical structure, synthesis, side effects, etc.,

- 9 -

of non-steroidal anti-inflammatory agents, reference may be had to standard texts, including Antiinflammatory and Anti-Rheumatic Drugs, K. D. Rainsford, Vol. I-III, CRC Press, Boca Raton, (1985), and Anti-inflammatory Agents, Chemistry and Pharmacology, 1, R. A. Scherrer, et al., Academic Press, New York (1974).

Specific non-steroidal anti-inflammatory agents useful in the compositions of the present invention include, but are not limited to:

- 1) the oxicams, such as piroxicam, isoxicam, tenoxicam, sudoxicam, and CP-14,304;
- 2) the salicylates, such as aspirin, disalcid, benorylate, trilisate, safapryn, solprin, diflunisal, and fendosal;
- 3) the acetic acid derivatives, such as diclofenac, fenclofenac, indomethacin, sulindac, tolmetin, isoxepac, furofenac, tiopinac, zidometacin, acematacin, fentiazac, zomepiract, clidanac, oxepinac, and felbinac;
- 4) the fenamates, such as mefenamic, meclofenamic, flufenamic, niflumic, and tolfenamic acids;
- 5) the propionic acid derivatives, such as ibuprofen, naproxen, benoxaprofen, flurbiprofen, ketoprofen, fenoprofen, fenbufen, indoprofen, pirprofen, carprofen, oxaprozin, pranoprofen, miroprofen, tioxaprofen, suprofen, alminoprofen, and tiaprofenic; and
- 6) the pyrazoles, such as phenylbutazone, oxyphenbutazone, feprazone, azapropazone, and trimethazone.

Mixtures of these non-steroidal anti-inflammatory agents may also be employed, as well as the pharmaceutically-acceptable salts and esters of these agents. For example, etofenamate, a flufenamic acid derivative, is particularly useful for topical application. Of the nonsteroidal anti-inflammatory agents, ibuprofen, naproxen, flufenamic acid, mefenamic acid, meclofenamic acid, piroxicam and felbinac are preferred; ibuprofen, naproxen, and flufenamic acid are most preferred.

Another class of anti-inflammatory agents which are useful in the present invention are the anti-inflammatory agents disclosed in

- 10 -

U.S. Patent No. 4,708,966, Loomans et al., issued November 24, 1987. This patent discloses a class of nonsteroidal anti-inflammatory compounds which comprise specifically substituted phenyl compounds, especially substituted 2,6-di-tert-butyl phenol derivatives. For example, compounds selected from 4-(4'-pentyn-3'-one)-2,6-di-t-butylphenol; 4-(5'-hexynoyl)-2,6-di-t-butylphenol; 4-((S)-(-)-3'-methyl-5'-hexynoyl)-2,6-di-t-butylphenol; 4-((R)-(+)-3'-methyl-5'-hexynoyl)-2,6-di-t-butylphenol; and 4-(3',3'-dimethoxypropionyl)-2,6-di-t-butylphenol are useful in the present invention.

Yet another class of anti-inflammatory agents which are useful in the present invention are those disclosed in U.S. Patent No. 4,912,248, Mueller, issued March 27, 1990. This patent discloses compounds and diastereomeric mixtures of specific 2-naphthyl-containing ester compounds, especially naproxen ester and naproxol ester compounds, having two or more chiral centers.

Finally, so-called "natural" anti-inflammatory agents are useful in the present invention. For example, candelilla wax, alpha bisabolol, aloe vera, Manjistha (extracted from plants in the genus Rubia, particularly Rubia Cordifolia), and Guggal (extracted from plants in the genus Commiphora, particularly Commiphora Mukul), may be used.

#### C. Anti-Oxidants/Radical Scavengers

In a preferred wrinkle and atrophy regulating composition of the present invention, an anti-oxidant/radical scavenger is included as an active agent along with the active compound. The inclusion of an anti-oxidant/radical scavenger increases the wrinkle regulating benefits of the composition.

A safe and effective amount of an anti-oxidant/radical scavenger may be added to the compositions of the present invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition.

Anti-oxidants/radical scavengers such as ascorbic acid (vitamin C) and its salts, tocopherol (vitamin E), tocopherol sorbate, other esters of tocopherol, butylated hydroxy benzoic acids and their

- 11 -

salts, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (commercially available under the tradename Trolox®), gallic acid and its alkyl esters, especially propyl gallate, uric acid and its salts and alkyl esters, sorbic acid and its salts, the ascorbyl esters of fatty acids, amines (e.g., N,N-diethylhydroxylamine, amino-guanidine), sulfhydryl compounds (e.g., glutathione), and dihydroxy fumaric acid and its salts may be used.

#### D. Chelators

In a preferred wrinkle and atrophy regulating composition of the present invention, a chelating agent is included as an active agent along with the active compound. As used herein, "chelating agent" means an active agent capable of removing a metal ion from a system by forming a complex so that the metal ion cannot readily participate in or catalyze chemical reactions. The inclusion of a chelating agent increases the wrinkle regulating benefits of the composition.

A safe and effective amount of a chelating agent may be added to the compositions of the present invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition. Chelators useful in compositions of the present invention are disclosed in U.S. Patent Application Serial No. 251,910,\* Bissett, Bush & Chatterjee, filed October 4, 1988, incorporated herein by reference. Preferred chelators useful in compositions of the present invention are furildioxime and derivatives thereof, more preferably amphi-2-furildioxime.

#### E. Retinoids

In a preferred wrinkle and atrophy regulating composition of the present invention, a retinoid, preferably retinoic acid, is included as an active agent along with the active compound. The inclusion of a retinoid increases the wrinkle regulating benefits of the composition. A safe and effective amount of a retinoid may be added to the compositions of the present invention, preferably from about 0.001% to about 2%, more preferably from about 0.01% to about 1% of the composition. As used herein, "retinoid" includes all natural and/or synthetic analogs of Vitamin A or retinol-like (\*Equivalent = EP Appln. 0313305, published 26 April 1989.)

- 12 -

compounds which possess the biological activity of Vitamin A in the skin as well as the geometric isomers and stereoisomers of these compounds, such as all-trans retinoic acid and 13-cis-retinoic acid.


F. Benzofuran Derivatives

5 In a preferred wrinkle and atrophy regulating composition of the present invention, a benzofuran derivative, preferably amiodarone, is included as an active agent along with the active compound. The inclusion of a benzofuran derivative increases the wrinkle regulating benefits of the composition.

10 A safe and effective amount of a benzofuran derivative may be added to the compositions of the present invention, preferably from about 0.01% to about 20%, more preferably from about 0.1% to about 10%, of the composition. Benzofuran derivatives useful in the present invention are disclosed in U.S. Patent Application Serial  
15 No. 674,628,\* Chatterjee and Kapoor, filed March 25, 1991, incorporated herein by reference.

G. N-acetyl-L-cysteine Derivatives

Also preferred for use herein are compounds having the structure



25 or a pharmaceutically-acceptable salt thereof.

$\text{R}^1$  is selected from the group consisting of nil and a  $\text{C}_1$ - $\text{C}_{18}$  alkyl, preferably  $\text{C}_1$ - $\text{C}_7$ , more preferably  $\text{C}_1$ - $\text{C}_3$ , more preferably still  $\text{C}_1$  alkyl.

$\text{R}^2$  is selected from the group consisting of nil, -H,  $\text{C}_1$ - $\text{C}_{18}$  alkyl and  $-\text{C}(=\text{O})-\text{R}^4$ ; preferably -H and,  $\text{C}_1$ - $\text{C}_{18}$  alkyl, more preferably -H. In one embodiment,  $\text{R}^2$  is preferably a  $\text{C}_1$ - $\text{C}_{18}$  alkyl, more preferably  $\text{C}_1$ - $\text{C}_7$ , more preferably  $\text{C}_1$ - $\text{C}_3$ , more preferably still  $\text{C}_1$ .

$\text{R}^3$  is selected from the group consisting of -H, and  $\text{C}_1$ - $\text{C}_{18}$  alkyl, preferably -H. In one embodiment,  $\text{R}^3$  is preferably a  $\text{C}_1$ - $\text{C}_{18}$

35 (\*Equivalent = WO92/07559, published 14 May 1992.)

- 13 -

alkyl, more preferably C<sub>1</sub>-C<sub>7</sub>, more preferably C<sub>1</sub>-C<sub>3</sub>, more preferably still C<sub>1</sub>.

R<sup>4</sup> is a C<sub>1</sub>-C<sub>18</sub> alkyl; preferably C<sub>1</sub>-C<sub>7</sub>; more preferably C<sub>1</sub>-C<sub>3</sub>; more preferably still C<sub>1</sub>.

5 In another embodiment, both R<sup>1</sup> and R<sup>2</sup> are nil and the carbonyl carbon and the sulfur adjacent R<sup>1</sup> and R<sup>2</sup>, respectively, are covalently bonded to form a cyclic ring. Otherwise, both R<sup>1</sup> and R<sup>2</sup> are other than nil.

10 Preferred pharmaceutically-acceptable salts of the active compound include, but are not limited to, sodium, potassium, magnesium, calcium, lithium, rubidium, strontium, aluminum, boron, silicon and zinc salts of the active compound.

15 Compositions of the present invention comprise from about 0.01% to about 50% of the active compound, preferably from about 0.1% to about 20%, more preferably from about 2% to about 5%.

Zinc complexes which may be formed by zinc and the active compound are useful in the compositions of the present invention.

#### H. Skin Protectants

20 In a preferred wrinkle and atrophy regulating composition of the present invention, a safe and effective amount of a skin protectant may be added to the compositions of the present invention, the skin protectant preferably comprises from about 0.001% to about 2%, more preferably from about 0.01% to about 1% of the composition. Useful skin protectants are disclosed in the  
25 Federal Register Vol. 48, No. 32 and include allantoin, aluminum hydroxide gel, bismuth subnitrate, boric acid, calamine, cocoa butter, corn starch, dimethicone, glycerin, kaolin, live yeast cell derivative, petrolatum, shark liver oil, sodium bicarbonate, sulfur, tannic acid, white petrolatum, zinc acetate, zinc carbonate and zinc  
30 oxide and mixtures thereof.

Other useful components include hormones such as pregnenolone and estrogens. Also useful are the alpha-, or beta-hydroxy acids or alpha-keto acids or derivatives thereof as disclosed in U. S. Patent 4,234,599 to Van Scott et al., issued November 18, 1980 which is  
35 incorporated by reference herein. Useful members of this class



- 14 -

include alpha-hydroxy-butyric acid, alpha-hydroxyisobutyric acid, alpha-hydroxyisocaproic acid, alpha-hydroxyisovaleric, atrolactic acid, beta-hydroxybutyric acid, beta-phenyl lactic acid, beta-phenylpyruvic acid, citric acid ethyl pyruvate, galacturonic acid, glucoheptonic acid, glucoheptono 1,4-lactone, gluconic acid, gluconolactone glucuronic acid, glucuronolactone, glycolic acid, isopropyl pyruvate, lactic acid, malic acid, amndelic acid, emthyl pyruvate, mucic acid, pyruvic acid, saccharic acid, saccharic acid 1,4-lactone, tartaric acid and tartronic acid.

10 I. Vitamins

Various vitamins may also be included in the compositions of the present invention. For example, Vitamin A, ascorbic acid, Vitamin B, biotin, panthothenic acid, Vitamin D, Vitamin E and mixtures thereof and derivatives thereof may be used.

15 Pharmaceutical Carriers

In a preferred embodiment, treatment will employ the use of a topical pharmaceutical composition comprising the active compound and a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier", as used herein, means one or more compatible solid or liquid filler diluents or microencapsulating substances which are suitable for administration to a human or lower animal. Pharmaceutically-acceptable carriers must be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the human or lower animal being treated. A safe and effective amount of carrier is from about 50% to about 99.99%, preferably from about 99.9% to about 80%, more preferably from about 98% to about 95%, of the composition.

Variations in formulation of these carriers will result in a wide variety of products which fall within the scope of the present invention.

The topical pharmaceutical compositions of the present invention may be made into a wide variety of product types. These include, but are not limited to solutions, lotions, creams, beach products, gels, sticks, sprays, pads, ointments, pastes, mousses and cosmetics. These product types may comprise several types of

- 15 -

carrier systems including, but not limited to solutions, emulsions, gels and solids.

The topical pharmaceutical compositions of the present invention formulated as solutions typically include a pharmaceuti-  
5 cally-acceptable aqueous or organic solvent. The terms "pharmaceutically-acceptable aqueous solvent" and "pharmaceutically-acceptable organic solvent" refer to a solvent which is capable of having dispersed or dissolved therein the active compound, and possesses acceptable safety properties (e.g., irritation and  
10 sensitization characteristics). Water is a typical aqueous solvent. Examples of suitable organic solvents include: propylene glycol, butylene glycol, polyethylene glycol (200-600), polypropylene glycol (425-2025), glycerol, 1,2,4-butanetriol, sorbitol esters, 1,2,6-hexanetriol, ethanol, isopropanol, butanediol, and mixtures  
15 thereof. Preferably, these solutions contain from about 0.01% to about 50% of the active compound, more preferably from about 0.1% to about 20%; and from about 1% to about 80% of an acceptable aqueous or organic solvent, more preferably from about 1% to about 40%.

If the topical pharmaceutical compositions of the present  
20 invention are formulated as an aerosol and applied to the skin as a spray-on, a propellant is added to a solution composition. A more complete disclosure of propellants useful herein can be found in Sagarin, Cosmetics Science and Technology, 2nd Edition, Vol. 2, pp. 443-465 (1972).

25 Topical pharmaceutical compositions of the present invention may be formulated as a solution comprising an emollient. An example of a composition formulated in this way would be a sunscreen-containing product. Preferably, such compositions contain from about 0.1% to about 50% of the active compound and from about  
30 2% to about 50% of a topical pharmaceutically-acceptable emollient.

As used herein, "emollients" refer to materials used for the prevention or relief of dryness, as well as for the protection of the skin. A wide variety of suitable emollients are known and may be used herein. Sagarin, Cosmetics Science and Technology, 2nd

- 16 -

Edition, Vol. 1, pp. 32-43 (1972), incorporated herein by reference, contains numerous examples of suitable materials.

5 A lotion can be made from a solution carrier system. Lotions preferably comprise from about 0.1% to about 20%, more preferably from about 1% to about 5%, of the active compound; from about 1% to about 20%, preferably from about 5% to about 10%, of an emollient; and from about 50% to about 90%, preferably from about 60% to about 80%, water.

10 Another type of product that may be formulated from a solution carrier system is a cream. A cream of the present invention would preferably comprise from about .1% to about 20%, more preferably from about 1% to about 5%, of the active compound; from about 5% to about 50%, preferably from about 10% to about 20%, of an emollient, and from about 45% to about 85%, preferably from about 50% to about 75%, water.

15 Yet another type of product that may be formulated from a solution carrier system is an ointment. An ointment may comprise a simple base of animal or vegetable oils or semi-solid hydrocarbons (oleaginous). Ointments may also comprise absorption ointment bases which absorb water to form emulsions. Ointment carriers may also be water soluble. An ointment may also comprise from about 2% to about 10% of an emollient plus from about 0.1% to about 2% of a thickening agent. A more complete disclosure of thickening agents useful herein can be found in Segarin, Cosmetics, Science and Technology, 20 2nd Edition, Vol. 1, pp. 72-73 (1972).

25 If the carrier is formulated as an emulsion, from about 1% to about 10%, preferably from about 2% to about 5%, of the carrier system comprises an emulsifier. Emulsifiers may be nonionic, anionic or cationic. Suitable emulsifiers are disclosed in, for example, U.S. Patent 3,755,560, issued August 28, 1973, Dickert et al.; U.S. Patent 4,421,769, issued December 20, 1983, Dixon et al.; and McCutcheon's Detergents and Emulsifiers, North American Edition, 30 pages 317-324 (1986); the disclosures of which are incorporated herein by reference. Preferred emulsifiers are anionic or nonionic, although the other types may also be used.

- 17 -

Lotions and creams can be formulated as emulsions as well as solutions. Preferably such lotions comprise from about 0.1% to about 20%, more preferably from about 1% to about 5%, of the active compound; from about 1% to about 20%, preferably from about 5% to about 10%, of an emollient; from about 25% to about 75%, preferably from about 45% to about 95%, water; and from about 0.1% to about 10%, preferably from about 0.5% to about 5%, of an emulsifier. Such creams would preferably comprise from about 0.1% to about 20%, more preferably from about 1% to about 5%, of the active compound; from about 1% to about 20%, preferably from about 5% to about 10%, of an emollient; from about 20% to about 80%, preferably from about 30% to about 70%, water; and from about 1% to about 10%, preferably from about 2% to about 5%, of an emulsifier.

Single emulsion skin care preparations, such as lotions and creams, of the oil-in-water type and water-in-oil type are well-known in the cosmetic art and are useful in the present invention. Multiphase emulsion compositions, such as the water-in-oil-in-water type, as disclosed in U.S. Patent No. 4,254,105, Fakuda et al., issued March 3, 1981, incorporated herein by reference, are also useful in the present invention. In general, such single or multiphase emulsions contain water, emollients and emulsifiers as essential ingredients.

Triple emulsion carrier systems comprising an oil-in-water-in-silicone fluid emulsion composition as disclosed in U.S. Patent No. 4,960,764, Figueroa, issued October 2, 1990, are also useful in the present invention. Preferably, this triple emulsion carrier system can be combined with from about 0.1% to about 20%, more preferably from about 1% to about 5%, of the active compound to yield the topical pharmaceutical composition of the present invention.

Another emulsion carrier system useful in the topical pharmaceutical compositions of the present invention is a micro-emulsion carrier system. Such a system comprises from about 9% to about 15% squalane; from about 25% to about 40% silicone oil; from about 8% to about 20% of a fatty alcohol; from about 15% to about 30% of

- 18 -

polyoxyethylene sorbitan mono-fatty acid (commercially available under the trade name Tweens) or other nonionics; and from about 7% to about 20% water. This carrier system is preferably combined with from about 1% to about 5% of the active compound.

5 If the topical pharmaceutical compositions of the present invention are formulated as a gel or a cosmetic stick, a suitable amount of a thickening agent, as disclosed supra, is added to a cream or lotion formulation.

10 The topical pharmaceutical compositions of the present invention may also be formulated as makeup products such as foundations.

The topical pharmaceutical compositions of the present invention may also be formulated as medicated pads. Suitable examples of these pads are fully disclosed in U.S. Patents 4,891,227  
15 and 4,891,228, to Thaman et al., both issued January 2, 1990 the disclosures of which are incorporated herein.

The topical pharmaceutical compositions of the present invention may contain, in addition to the aforementioned components, a wide variety of additional oil-soluble materials and/or  
20 water-soluble materials conventionally used in topical compositions, at their art-established levels.

Various water-soluble materials may also be present in the compositions of this invention. These include humectants, proteins and polypeptides, preservatives and an alkaline agent. In addition,  
25 the topical compositions herein can contain conventional cosmetic adjuvants, such as dyes, opacifiers (e.g., titanium dioxide), pigments and perfumes.

The topical pharmaceutical compositions of the present invention may also include a safe and effective amount of a penetration enhancing agent. A preferred amount of penetration  
30 enhancing agent is from about 1% to about 5% of the composition. Another useful penetration enhancer for the present invention is the non-ionic polymer under the CTFA designation: polyacrylamide and isoparaffin and laureth-7, available as Sepigel from Seppic Corporation. Also useful is polyquaternium-32 and mineral oil known  
35

- 19 -

as SalCare SC92 available from Allied Colloids, Suffolk, VA. This is a class of cationic polymers which are generally described in U.S. Patent 4,628,078 to Glover et al. issued December 9, 1986 and U.S. Patent 4,599,379 to Flesher et al. issued July 8, 1986 both of which are incorporated by reference herein.

Examples of useful penetration enhancers, among others, are disclosed in U.S. Patents 4,537,776, Cooper, issued August 27, 1985; 4,552,872, Cooper et al., issued November 12, 1985; 4,557,934, Cooper, issued December 10, 1985; 4,130,667, Smith, issued December 19, 1978; 3,989,816, Rhaadhyaksha, issued November 2, 1976; 4,017,641, DiGiulio, issued April 12, 1977; and European Patent Application 0043738, Cooper et al., published January 13, 1982.

Other conventional skin care product additives may also be included in the compositions of the present invention. For example, collagen, hyaluronic acid, elastin, hydrolysates, primrose oil, jojoba oil, epidermal growth factor, soybean saponins, mucopolysaccharides, and mixtures thereof may be used.

#### Cleaning Compositions

The skin cleaning compositions of the present invention comprise, in addition to the active compound, a cosmetically-acceptable surfactant. The term "cosmetically-acceptable surfactant" refers to a surfactant which is not only an effective skin cleanser, but also can be used without undue toxicity, irritation, allergic response, and the like. Furthermore, the surfactant must be capable of being commingled with the active compound in a manner such that there is no interaction which would substantially reduce the efficacy of the composition for regulating skin wrinkles and/or skin atrophy.

The skin cleaning compositions of the present invention preferably contain from about 0.1% to about 20%, preferably from about 1% to about 5%, of the active compound and from about 1% to about 90%, more preferably from about 1% to about 10%, of a cosmetically-acceptable surfactant.

- 20 -

The physical form of the skin cleansing compositions is not critical. The compositions can be, for example, formulated as toilet bars, liquids, pastes, mousses, or pads.

5 The surfactant component of the compositions of the present invention are selected from anionic, nonionic, zwitterionic, amphoteric and ampholytic surfactants, as well as mixtures of these surfactants. Such surfactants are well-known to those skilled in the detergency art.

10 The cleaning compositions of the present invention can optionally contain, at their art-established levels, materials which are conventionally used in skin cleansing compositions.

Regulating Wrinkles and/or Skin Atrophy in Mammalian Skin

15 The amount of active components and frequency of treatment will vary widely depending upon the level of wrinkling and/or skin atrophy already in existence in the subject, the rate of further wrinkle formation and/or atrophy, and the level of regulation desired.

20 Preferably the composition is applied to the skin is via chronic topical application of a safe and effective amount of the composition to regulate wrinkles and/or atrophy in mammalian skin. The amount of actives and frequency of topical application to the skin can vary widely, depending upon personal needs, but it is suggested as an example that topical application range from about  
25 once per week to about 10 times daily, preferably from about twice per week to about 4 times daily, more preferably from about 3 times a week to about 3 times daily, most preferably about once or twice per day. The composition for topical application will comprise from about 0.01% to about 50%, preferably from about 0.1% to about 20%, more preferably from about 1% to about 5% of the active compound.  
30 By "chronic" application, it is meant herein that the period of topical application may be over the lifetime of the subject, preferably for a period of at least about three weeks, more preferably from about three months to about twenty years, more preferably from about six months to about ten years, more preferably

35

- 21 -

still from about one year to about five years, thereby resulting in regulation of wrinkles and/or atrophy in mammalian skin.

The compositions of the present involves applying both a safe and effective amount of the active compound and a safe and effective amount of one or more of a sunscreensing agent, anti-inflammatory agent, vitamin, anti-oxidant/radical scavenging agent, chelating agent, retinoid, N-acetyl-L-cysteine derivative, skin protectant and/or benzofuran derivative to the skin simultaneously. As used herein, "simultaneous application" or "simultaneously" means applying the agents to the skin at the same situs on the body at about the same time. Though this can be accomplished by applying the agents separately to the skin, preferably a composition comprising all the desired agents commingled is applied to the skin. The amount of sunscreensing agent applied is generally from about 0.02 mg to about 1.0 mg per  $\text{cm}^2$  skin. The amount of anti-inflammatory agent applied is generally from about 0.005 mg to about 0.5 mg, preferably from about 0.01 mg to about 0.1 mg per  $\text{cm}^2$  skin. The amount of anti-oxidant/radical scavenging agent generally applied is from about 0.001 mg to about 1.0 mg, preferably from about 0.05 mg to about 0.5 mg per  $\text{cm}^2$  skin. The amount of chelating agent generally applied is from about .001 mg to about 1.0 mg, preferably from about 0.01 mg to about 0.5 mg, more preferably from about 0.05 mg to about 0.1 mg per  $\text{cm}^2$  skin. The amount of retinoid applied is generally from about 0.00001 mg to about 0.02 mg per  $\text{cm}^2$  skin, preferably from about 0.001 mg to about 0.01 mg per  $\text{cm}^2$  skin. The amount of benzofuran derivative applied is generally from about 0.001 mg to about 1.0 mg/ $\text{cm}^2$  skin per application, preferably from about 0.01 to about 0.5 mg/ $\text{cm}^2$  skin per application. The amount of active compound applied is generally from about 0.001 mg to about 1.0 mg per  $\text{cm}^2$  skin per application, preferably from about 0.01 mg to about 0.5 mg per  $\text{cm}^2$ , more preferably from about 0.05 to about 0.25 mg/ $\text{cm}^2$  skin per application.

The following examples further describe and demonstrate the preferred embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration, and



- 22 -

are not to be construed as limitations of the present invention since many variations thereof are possible without departing from its spirit and scope.

EXAMPLE I

A pad of the present invention is made as follows:

<u>Pad Composition</u>	<u>Weight %</u>
<u>Substrate A</u>	
Cellulose-based nonwoven <sup>1</sup>	100.0

<u>Substrate B</u>	
Polyester (denier = 6) <sup>2</sup>	45.0
Orlon (denier = 8) <sup>3</sup>	15.0
Styrene-butadiene resin <sup>4</sup>	40.0

<u>Laminate</u>	
Polyethylene Powder Melt <sup>5</sup>	100.0

<u>Active Composition</u>	<u>Weight %</u>
Salicylic acid	2.0
Na Methyl cocoyl taurate	3.0
C <sub>2</sub> H <sub>5</sub> OH (95% ethanol)	35.0
Witch Hazel distillate	5.0
Quaternium-22	1.0
Menthol	0.1
Aloe Vera Gel	0.5
Fragrance	0.05
Water	q.s.

<sup>1</sup> Obtained from James River as Airtex Spec 382.

<sup>2</sup> Obtained from Eastern Chemical Company.

<sup>3</sup> Obtained from American Cyanamid.

<sup>4</sup> Obtained from Reichold as tylac 68-500 (ratio of styrene to butadiene 80:20).

<sup>5</sup> Obtained from Quantum Chemical as microthene powder.

- 23 -

Substrate A has a basis weight of about 55 grams per square yard and a loft of about 35 mills. Substrate B has a basis weight of about 65 grams per square yard and a loft of about 70 to 80 mills. The two materials are laminated together by applying a thin coat of Polyethylene power to Substrate A and heating with IR lamps. Substrate A and B are then joined at a hip roll to compress and bond the materials. The resulting nonwoven fabric has a loft of about 90 to 100 mills. The resulting material is then cut into an oval shape (5 cm. x 7 cm.). The active components are combined to form a solution and the pad composition is saturated in this solution.

This composition is useful for topical application to regulate skin wrinkles and/or skin atrophy. Use of an amount of the composition to deposit about 2 mg/cm<sup>2</sup> of the active compound to the skin is appropriate.

#### EXAMPLE II

A pad of the present invention is made by combining the following components as in Example I:

<u>Pad Composition</u>	<u>Weight %</u>
<u>Substrate A</u>	
Cellulose-based nonwoven	100.0
<u>Substrate B</u>	
Polyester (denier = 6)	45.0
Orlon (denier = 8)	15.0
Styrene-butadiene resin	40.0
<u>Laminate</u>	
Polyethylene Powder Melt	100.0

- 24 -

	<u>Active Composition</u>	<u>Weight %</u>
	Salicylic acid	2.0
	C <sub>2</sub> H <sub>5</sub> OH (95% ethanol)	35.0
	Glycerin	2.0
5	Aloe Vera Gel	1.0
	Menthol	0.05
	Triethanol Amine	0.7
	Fragrance	0.15
	Water	q.s.

10

This composition is useful for topical application to regulate skin wrinkles and/or skin atrophy. Use of an amount of the composition to deposit about 2 mg/cm<sup>2</sup> of the active compound to the skin is appropriate.

15

### EXAMPLE III

A topical composition is prepared by combining the following components utilizing conventional mixing techniques.

	<u>Active Composition</u>	<u>Weight %</u>
20	Salicylic acid	1.25
	Ascorbic acid	5.00
	Na Methyl cocoyl taurate	1.5
	C <sub>2</sub> H <sub>5</sub> OH (95% ethanol)	45.0
	Witch Hazel distillate	5.0
25	Quaternium-22	1.0
	Menthol	0.05
	Fragrance	0.05
	Water	41.15

30 This composition is useful for topical application to regulate skin wrinkles and/or skin atrophy. Use of an amount of the composition to deposit about 2 mg/cm<sup>2</sup> of the active compound to the skin is appropriate.

35

- 25 -

EXAMPLE IV

A topical composition is made by combining the following components using conventional mixing technology.

	<u>Ingredient</u>	<u>W/W%</u>
5	Water, Purified	54.0
	Alcohol SD 40	40.0
	Polyacrylamide and C <sub>13-14</sub>	
	Isoparaffin and Laureth-7 <sup>1</sup>	4.0
	Salicylic Acid	2.0

10 <sup>1</sup>Available as Sepigel from Seppic Corporation.

Water is added to a suitable size container. While mixing at a moderate speed (300 rpm), the polyacrylamide isoparaffin and laureth-7 is added to the water. Separately, the alcohol is placed in a container and covered. Using a Lightnin' Mixer with a 3 blade paddle prop, the salicylic acid is added to the alcohol and mixed at 15 a low speed (100 rpm) until all salicylic acid is dissolved. The alcohol is slowly added to the water phase to form a gel. The resulting gel is mixed at moderate speed until uniform.

This composition is useful for topical application to regulate 20 skin wrinkles and/or skin atrophy. Use of an amount of the composition to deposit about 2 mg/cm<sup>2</sup> of the active compound to the skin is appropriate.

25

30

35

- 26 -

EXAMPLE V

A topical composition is made by combining the following components using conventional mixing technology as in Example IV.

	<u>Ingredient</u>	<u>W/W%</u>
5	Water	q.s.
	Alcohol SD 40	40.0
	Salcare SC92 <sup>1</sup>	3.0
	Salicylic Acid	2.0
	Menthol	0.05
10	Ng <sub>2</sub> EDTA	0.05
	Glycerin	2.00

<sup>1</sup>Salcare SC92 is a copolymer of acylamide and a cationic acrylate available from Allied Colloids.

15 This composition is useful for topical application to regulate skin wrinkles and/or skin atrophy. Use of an amount of the composition to deposit about 2 mg/cm<sup>2</sup> of the active compound to the skin is appropriate.

20

25

30

35

- 27 -

## CLAIMS:

1. A composition for regulating wrinkles or atrophy in mammalian skin comprising:
  - (a) a safe and effective amount of salicylic acid;
  - (b) another active agent selected from the group consisting of a safe and effective amount of a sunscreen, an anti-inflammatory agent, a vitamin, an anti-oxidant, a chelator, a retinoid, a benzofuran derivative, an N-acetyl-L-cysteine derivative and a skin protectant, derivatives thereof and mixtures thereof; and
  - (c) a pharmaceutically-acceptable carrier.
2. The composition of Claim 1 wherein the composition comprises from 0.01% to 50% of salicylic acid, preferably from 0.1% to 20% of salicylic acid.
3. The composition of Claim 2 wherein the pharmaceutically-acceptable carrier is a topical carrier.
4. The composition of Claim 3 wherein the topical carrier comprises:
  - (a) from 10 to 60 weight percent of  $C_2H_5OH$  or  $C_3H_7OH$ ;
  - (b) from 30 to 80 weight percent of water; and
  - (c) from 0.2 to 5.0 weight percent of sodium methyl cocoyl taurate or sodium methyl oleoyl taurate; the composition having a pH value of from 2 to 3.5.
5. The composition of Claim 4 wherein the composition additionally comprises a safe and effective amount of a sunscreensing agent.
6. The composition of Claim 5 wherein said sunscreen is selected from the group consisting of 2-ethylhexyl-p-methoxycinnamate,

- 28 -

butylmethoxydibenzoylmethane, 2-hydroxy-4-methoxybenzophenone, octyldimethyl-p-aminobenzoic acid and mixtures thereof.

7. The composition of Claim 4 wherein said anti-inflammatory agent is selected from the group consisting of hydrocortisone, hydroxyltriamcinolone, alpha-methyl dexamethasone, dexamethasone-phosphate, beclomethasone dipropionate, clobetasol valerate, desonide, desoxymethasone, desoxycorticosterone acetate, dexamethasone, dichlorisone, diflorasone diacetate, difluocortolone valerate, fluadrenolone, flucolorolone acetonide, fludrocortisone, flumethasone pivalate, fluosinolone acetonide, fluocinonide, flucortine butylester, flucortolone, fluprednidene (fluprednylidene) acetate, flurandrenolone, halcinonide, hydrocortisone acetate, hydrocortisone butyrate, methylprednisolone, triamcinolone acetonide, cortisone, cortodoxone, flucetonide, fludrocortisone, difluorosone diacetate, fluradrenolone acetonide, medrysone, amcinafel, amcinafide, betamethasone and the balance of its esters, chloroprednisone, chlorprednisone acetate, clocortelone, clescinalone, dichlorisone, difluprednate, flucoloronide, flunisolide, fluoromethalone, fluperolone, fluprednisolone, hydrocortisone valerate, hydrocortisone cyclopentylpropionate, hydrocortamate, meprednisone, paramethasone, prednisolone, prednisone, beclomethasone dipropionate, triamcinolone, piroxicam, isoxicam, tenoxicam, sudoxicam, CP-14,304, aspirin, disalcid, benorylate, trilisate, safapryn, solprin, diflunisal, and fendosal, diclofenac, fenclofenac, indomethacin, sulindac, tolmetin, isoxepac, furofenac, tiopinac, zidometacin, acematacin, fentiazac, zomepiract, clidanac, oxepinac, and felbinac, mefenamic, meclofenamic, flufenamic, niflumic, and tolfenamic acid, ibuprofen, naproxen, benoxaprofen, flurbiprofen, ketoprofen, fenoprofen, fenbufen, indoprofen, pirprofen, carprofen, oxaprozin, pranoprofen, miroprofen, tioxaprofen, suprofen, alminoprofen, and tiaprofenic,

- 29 -

phenybutazone, oxyphenbutazone, feprazone, azapropazone, and trimethazone and mixtures thereof.

- 5 8. The composition of Claim 4 wherein said anti-oxidant or radical scavenger is selected from the group consisting of ascorbic acid, tocopherol, butylated hydroxy benzoic acids, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid, gallic acid, uric acid, sorbic acid, N,N-diethylhydroxylamine, amino-guanidine), sulfhydryl compounds, and dihydroxy fumaric  
10 acid, derivatives thereof and mixtures thereof.
9. The composition of Claim 4 wherein said retinoid is selected from the group consisting of all-trans retinoic acid and 13-cis-retinoic acid and mixtures thereof.  
15
10. The composition of Claim 4 wherein said skin protectant is selected from the group consisting of allantoin, aluminum hydroxide gel, bismuth subnitrate, boric acid, calamine, cocoa butter, corn starch, dimethicone, glycerin, kaolin, live yeast  
20 cell derivative, petrolatum, shark liver oil, sodium bicarbonate, sulfur, tannic acid, white petrolatum, zinc acetate, zinc carbonate and zinc oxide and mixtures thereof.



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 92/09737

**I. CLASSIFICATION OF SUBJECT MATTER** (If several classification symbols apply, indicate all)\*

According to International Patent Classification (IPC) or to both National Classification and IPC

Int.Cl. 5 A61K7/48

**II. FIELDS SEARCHED**Minimum Documentation Searched<sup>7</sup>

Classification System	Classification Symbols
Int.Cl. 5	A61K

Documentation Searched other than Minimum Documentation  
to the extent that such Documents are included in the Fields Searched<sup>8</sup>**III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup>**

Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
X	US,A,5 017 367 (R. G. STOJKOSKI) 21 May 1991 see column 3, line 57 - line 61 see column 3, line 40 - line 47; claims 1-3; example 1 ---	1-3,10
X	JAPANESE PATENTS GAZETTE Week 8322, Derwent Publications Ltd., London, GB; & JP,A,55 040 632 (NIPPON SOLID CO LTD) 16 September 1978 see abstract ---	1
X	EP,A,0 434 628 (P. HERZOG) 26 June 1991 see page 6, line 1 - line 4; examples 1-3 ---	1-3,8,9
	---	-/--

<sup>10</sup> Special categories of cited documents:<sup>"A"</sup> document defining the general state of the art which is not considered to be of particular relevance<sup>"E"</sup> earlier document but published on or after the international filing date<sup>"L"</sup> document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)<sup>"O"</sup> document referring to an oral disclosure, use, exhibition or other means<sup>"P"</sup> document published prior to the international filing date but later than the priority date claimed<sup>"T"</sup> later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention<sup>"X"</sup> document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step<sup>"Y"</sup> document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.<sup>"A"</sup> document member of the same patent family**IV. CERTIFICATION**

Date of the Actual Completion of the International Search

27 JANUARY 1993

Date of Mailing of this International Search Report

16.02.93

International Searching Authority

EUR PEAN PATENT OFFICE

Signature of Authorized Officer

SIERRA GONZALEZ

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	US,A,4 608 370 (R. B. ARONSOHN) 26 August 1986 see column 3, line 26; example 1; tables I,II ---	1-3
X	US,A,3 821 370 (L. T. TENTA) 28 June 1974 see column 2, line 3 - line 26; example 1 ---	1-3,8,10
X	EP,A,0 378 936 (L'OREAL) 25 July 1990 see the whole document -----	1-3,5,9

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9209737  
SA 66954

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

27/01/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5017367	21-05-91	None	
EP-A-0434628	26-06-91	CH-A- 679981	29-05-92
US-A-4608370	26-08-86	None	
US-A-3821370	28-06-74	US-A- 3949072	06-04-76
EP-A-0378936	25-07-90	LU-A- 87408	10-07-90
		CA-A- 2005695	16-06-90
		JP-A- 2218602	31-08-90